

CORRECTION

Open Access



Correction: The role of the KEAP1-NRF2 signaling pathway in form deprivation myopia guinea pigs

Zhiming Gu^{1,2†}, Jiayu Meng^{3†}, Weiqi Zhong^{1,2}, Changjun Lan^{1,2}, Qingqing Tan^{1,2}, Xiaoling Xiang^{1,2}, Hong Zhou^{1,2} and Xuan Liao^{1,2*}

Correction: *BMC Ophthalmol* 24, 497 (2024)
<https://doi.org/10.1186/s12886-024-03754-6>

Following publication of this article, it was brought to our attention that in Table 1 the reverse primer of NRF2 was incorrectly given as 5'-TTTGCAAGCATCCTCGTTC C-3'.

The correct NRF2 reverse primer should be 5'-TGTCCT GCTGCATGTTGTCT-3'.

The correct version of Table 1 can be found below:

Table 1 The mRNA sequences

Gene	Forward primer	Reverse primer
KEAP1	5'-ATGAGTGTACCCCGAAACCG-3'	5'-TACAGCAGGCGATT-GAGCAC-3'
NRF2	5'-GGCAACTACTCCAGGTTTCT-3'	5'-TGTCCTGCTGCAT-GTTGTCT-3'
SOD	5'-TGAGCCCTAATGGTGGTGA-3'	5'-TTTGCAAGCATCCTC-GTTCC-3'
GAPDH	5'-GGTATTCCTTCTCCCGTGC-3'	5'-CCAAATCC-GTTCCTCCGA-3'

The original article has been corrected.

The authors would like to apologize for any inconvenience caused to the readers.

Published online: 17 March 2025

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

[†]Zhiming Gu and Jiayu Meng contributed equally to this work.

The online version of the original article can be found at <https://doi.org/10.1186/s12886-024-03754-6>.

*Correspondence:

Xuan Liao
aleexand@163.com

¹Ophthalmology Department of Affiliated Hospital of North Sichuan Medical College, Medical School of Ophthalmology & Optometry, Nanchong, Sichuan Province 637000, China

²Medical School of Ophthalmology & Optometry, North Sichuan Medical College, Nanchong, Sichuan Province 637000, China

³Sichuan Provincial Key Laboratory for Human Disease Gene Study, Sichuan Provincial People's Hospital, University of Electronic Science and Technology of China, Chengdu 611731, China

